

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/341425571>

Distinct psychological profiles among college students with substance use: A cluster analytic approach

Article in *Addictive Behaviors* · May 2020

DOI: 10.1016/j.addbeh.2020.106477

CITATIONS

0

READS

78

7 authors, including:



Séverine Lannoy
Stanford University

30 PUBLICATIONS 167 CITATIONS

[SEE PROFILE](#)



Jessica Mange
Université de Caen Normandie

25 PUBLICATIONS 86 CITATIONS

[SEE PROFILE](#)



Pascale Leconte
Laboratoire COMETE INSERM 1075

25 PUBLICATIONS 213 CITATIONS

[SEE PROFILE](#)



Ludivine Ritz
Université de Caen Normandie

43 PUBLICATIONS 154 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Socio-emotional changes in brain-damaged patients [View project](#)



Brain mechanisms underlying episodic simulation of future sustainable behaviors [View project](#)



Distinct psychological profiles among college students with substance use: A cluster analytic approach

Séverine Lannoy^{a,b,c,*}, Jessica Mange^d, Pascale Leconte^e, Ludivine Ritz^d, Fabien Gierski^{a,f}, Pierre Maurage^b, Hélène Beaunieux^d

^a Université de Reims Champagne-Ardenne, URCA, C2S (EA6291), 57 rue Pierre Taittinger, 51571 Reims, France

^b Université catholique de Louvain, UCLouvain, Louvain Experimental Psychopathology Research Group (LEP), 10 Place Cardinal Mercier, 1348 Louvain-la-Neuve, Belgium

^c Stanford University, Department of Psychiatry and Behavioral Sciences, Stanford, CA, USA

^d Université de Caen-Normandie, UNICAEN, LPCN, Esplanade de la Paix, 14000 Caen, France

^e Université de Caen-Normandie, UMR-S 1075 INSERM/UNICAEN COMETE, 2 rue des Rochambelles, 14032 Caen, France

^f Groupe de Recherche sur l'Alcool et les Pharmacodépendances, GRAP, INSERM U1247, Chemin du Thil, 80025 Amiens, France

HIGHLIGHTS

- Substance use in young people is an important public health concern.
- Research has explored the psychological factors related to substance use.
- Few studies have considered multiple substance use and possible heterogeneity.
- We performed a cluster analysis based on impulsivity traits and consumption motives.
- Four subgroups were found, described by distinct profiles and substance use habits.

ARTICLE INFO

Keywords:
Motives
Impulsivity
College students
Cluster analysis

ABSTRACT

Substance use in youth is a central public health concern, related to deleterious consequences at psychological, social, and cognitive/cerebral levels. Previous research has identified impulsivity and consumption motives as key factors in the emergence of excessive substance use among college students. However, most studies have focused on a specific substance and have considered this population as a unitary group, ignoring the potential heterogeneity in psychological profiles. We used a cluster analytic approach to explore the heterogeneity in a large sample ($N = 2741$) of substance users (i.e., tobacco, alcohol, cannabis, ecstasy, cocaine, heroin) on impulsivity and consumption motives. We identified four clusters: The first two clusters, associated with good self-esteem, low anxiety, and moderate substance use, were respectively characterized by low impulsivity and consumption motives (Cluster 1) and by high social and enhancement motives without marked impulsivity (Cluster 2). The two other clusters were conversely related to low self-esteem and high anxiety, and characterized by high consumption motives (particularly conformity) together with elevated urgency (Cluster 3) and by globally increased impulsivity and consumption motives (Cluster 4). These two clusters were also associated with higher substance use. These results highlight the existence of distinct psychological profiles of substance users and underline the need to develop targeted prevention and intervention programs (e.g., focusing on the specific impulsivity facets and consumption motives presented by each subgroup). Based on these findings, we also suggest extending the exploration of distinct profiles of substance users by targeting other psychological variables (e.g., self-esteem).

* Corresponding author at: Stanford University, Department of Psychiatry and Behavioral Sciences, 401 Quarry Rd, Stanford, CA 94305, USA.

E-mail addresses: severine.lannoy.phd@gmail.com (S. Lannoy), jessica.mange@unicaen.fr (J. Mange), pascale.leconte@unicaen.fr (P. Leconte), ludivine.ritz@unicaen.fr (L. Ritz), fabien.gierski@univ-reims.fr (F. Gierski), pierre.maurage@uclouvain.be (P. Maurage), helene.beaunieux@unicaen.fr (H. Beaunieux).

<https://doi.org/10.1016/j.addbeh.2020.106477>

Received 30 December 2019; Received in revised form 13 May 2020; Accepted 15 May 2020

Available online 21 May 2020

0306-4603/ Published by Elsevier Ltd.

1. Introduction

Substance use is a major concern in Western societies and widespread among college students, as about 60% of young adults drink alcohol frequently, 10% are regular smokers, and more than 20% consume illicit drugs such as marijuana, cocaine, or heroin (Center for Behavioral Health Statistics and Quality, 2015). Moreover, substance use in young people is often associated with important consequences; such as poor academic performance, legal problems, interpersonal conflicts, suicidal ideations, or cognitive impairments (Ayala, Roseman, Winseman, & Mason, 2017; see Skidmore, Kaufman, & Crowell, 2016 for a review). The identification of people at-risk to consume such substances is thus crucial to further develop individualized prophylactic interventions (e.g., Grant, Stewart, O'Connor, Blackwell, & Conrod, 2007; Skidmore et al., 2016). Nevertheless, whereas previous works have identified the main psychological variables involved in the emergence and maintenance of substance use, it remains unclear whether most users show an identical psychological profile or whether subgroups with dissociated psychological characteristics exist.

The current literature related to the determinants of substance use in youth has identified the effect of common factors (e.g., peer influence, externalized behaviors), whereas the specific variables associated with each substance use appear more related to individual factors (e.g., gender, school years) (Fitzgerald, Mac Giollabhui, Dolphin, Whelan, & Dooley, 2018). Beyond these general variables, psychological studies have identified both trait and behavioral impulsivity as a critical factor influencing substance use, specifically the inability to take the expected consequences into consideration (Charles et al., 2016; Magallón-Neri, Díaz, Forns, Goti, & Castro-Fornieles, 2015) and the tendency to seek for new and enjoyable sensations (Charles et al., 2016; Malmberg et al., 2012). Importantly, trait impulsivity is considered to be a good predictor of treatment outcomes in substance use disorders (see Loree, Lundahl, & Ledgerwood, 2015 for a review), which underlines its involvement in the inability to control substance use and the need to understand its role in the consumption patterns observed among young people. Several conceptualizations of impulsivity have emerged in the last decades. Among them, the UPPS (Whiteside & Lynam, 2001) constitutes an internationally recognized model in addiction research. This model describes impulsivity through four facets reflecting both trait and behavioral components: Urgency (impetuous actions following intense emotional states), lack of Premeditation (absence of consequences consideration), lack of Perseverance (difficulty to stay focused on a demanding/boring task), and Sensation seeking (search for stimulations and new experiences). This model has allowed for the specification of impulsivity facets related to different types of substance use. Initial studies have revealed that impulsivity is generally associated with hazardous alcohol use (Shin, Hong, & Jeon, 2012), but more recent works have shown that, in a sample of youth with substance use disorders, high sensation seeking and lack of perseverance were prevalent among individuals with problematic alcohol use, urgency was related to cannabis consumption, and lack of perseverance was related to use of other drugs (Thomsen et al., 2018).

In addition, some studies have emphasized the importance of targeting subjective reasons given by young people for using substances (Dow & Kelly, 2013). From this perspective, a reliable way to evaluate consumption motives can be derived from Cooper (1994) motivational model (e.g., Hides et al., 2008; Patrick, Fairlie, & Lee, 2018). This model is among the most influential in addiction research and postulates that individuals consume drugs for positive and/or negative reinforcements related to internal and/or external sources. Accordingly, Cooper (1994) defines four motives, namely social order (positive-external), enhancement (positive-internal), conformity (negative-external), and coping (negative-internal). Studies have supported this model by showing the involvement of motives in substance use disorders (see Cooper, Kuntsche, Levitt, Barber, & Wolf, 2016 for a review; Lee, Derefinko, Davis, Milich, & Lynam, 2017). Indeed, whereas

substance users reported coping and social motives (Hides et al., 2008; Wong et al., 2013), enhancement and conformity did not seem specifically related to drug consumption (Hides et al., 2008). When exploring the comorbid use of alcohol and cannabis in youth, social motives were mainly related to alcohol use; enhancement motives predicted both alcohol and cannabis consumption, whereas coping motives were related to cannabis but not alcohol use (Skalisky, Wielgus, Aldrich, & Mezulis, 2019).

The joint influence of impulsivity and consumption motives has also been reported to predict cannabis (e.g., Hecimovic, Barrett, Darredeau, & Stewart, 2014) or alcohol (e.g., Jones, Chrissyanthakis, & Groom, 2014) use among young people. In particular, it has been shown that the associations between problematic drinking and sensation seeking or lack of premeditation were mediated by enhancement motives, and that the association with negative urgency was influenced by coping, social, and enhancement motives (Adams, Kaiser, Lynam, Charnigo, & Milich, 2012; Jones et al., 2014). However, possible interactions between these variables have to be further explored in multiple substance use. Indeed, although the current literature has led to a better understanding of the impulsivity facets and motives involved in different consumptions (e.g., Hides et al., 2008; Thomsen et al., 2018), there is a need for studies considering multiple substance use, which is a very common pattern in college students. Furthermore, research increasingly shows that it is central to consider the possible existence of subgroups characterized by distinct psychological profiles (e.g., Malmberg et al., 2012). Nevertheless, previous studies focusing on substance use have mainly explained heterogeneity through social and environmental variables (e.g., Assanangkornchai, Li, McNeil, & Saingam, 2018; Evans-Polce, Lanza, & Maggs, 2016; Schilling et al., 2017), without considering the role of central psychological factors. Heterogeneity of psychological profiles has been confirmed in young adults with excessive alcohol or cannabis use (e.g., Gierski et al., 2017; Lannoy, Billieux, Poncin, & Maurage, 2017; Martinez-Loredo, Fernandez-Hermida, De La Torre-Luque, & Fernandez-Artamendi, 2018; Pedersen, Thomsen, Pedersen, & Hesse, 2017), but the joint role of impulsivity facets and consumption motives in substance use remains to be established.

The current study aimed to explore the use of various substances in a sample of college students and particularly examined (1) the psychological profiles of individuals with substance use, and (2) the comorbidities and severity of substance use behaviors in each profile. Based on impulsivity and consumption motives, we used a cluster analytic approach to explore the existence of distinct psychological profiles among college students. We then evaluated the differences between clusters regarding the prevalence and intensity of substance use, the presence of anxiety, and self-esteem level. Indeed, high anxiety and low self-esteem are crucial factors associated with substance use in young people (e.g., Blank, Connor, Gray, & Tustin, 2016; Walther, Morgenstern, & Hanewinkel, 2012) and may help to characterize the profiles of substance users. Capitalizing on European and French prevalences, we focused this study on several substances, namely tobacco, alcohol, cannabis, ecstasy, cocaine, and heroin (European Monitoring Center for Drugs and Drug Addiction¹, 2015). According to previous studies (e.g., Lammers, Kuntsche, Engels, Wiers, & Kleinjan, 2013; Lannoy et al., 2017), we hypothesized that profiles combining high impulsivity and stronger motives would be related to increased and harmful substance use.

2. Materials and methods

2.1. Participants and design

We recruited a convenient sample of 2741 college students from the *University of Caen Normandie* (France; academic year 2016–2017). This

¹ Observatoire Européen des Drogues et des Toxicomanie.




Fig. 1. Flowchart of data processing and inclusion.

study was part of a large research project exploring substance consumption in young adults (ADUC; Alcohol and Drug at University of Caen). Participants were contacted by email to answer an online survey screening substance use in university students (anonymity was guaranteed). A total of 34,215 emails were sent and 2741 participants were included in the study after removing missing data and abstainers (Fig. 1). There was no other inclusion or exclusion criteria. The current response rate (13.87%²) as well as the ratio between completed responses and included participants (62.84%) were comparable to previous studies among college students (e.g., Ehret, Ghaidarov, & LaBrie, 2013; McCabe, Boyd, Couper, Crawford, & D'arcy, 2002; Neighbors, Dillard, Lewis, Bergstrom, & Neil, 2006). The study protocol was approved by the National computer and freedom commission (CNIL³; file number u24-20171109-01R1) and conducted according to the Declaration of Helsinki. All students provided informed consent before taking part in the study.

2.2. Measures

The online survey was implemented through LimeSurvey software and assessed sociodemographic (i.e., age, gender, mother tongue) and academic (i.e., attendance, study years, diploma, grade) variables, substance use (tobacco, alcohol, cannabis, ecstasy, cocaine, heroin; see

Supplementary Table for reliability coefficients), motives in association with substance use, impulsivity (UPPS-P; Billieux et al., 2012), and trait anxiety (State-Trait Anxiety Inventory, STAI; Bruchon-Schweitzer & Paulhan, 1993). Also, we assessed self-esteem through a unique item (i.e., “I have a good self-esteem”), on a Likert scale from 0 “not at all” to 5 “absolutely” (Robins, Hendin, & Trzesniewski, 2001). Regarding consumption motives, we used the short form questionnaire developed by Kuntsche and Kuntsche (2009) for all substances (i.e. “if you think about all the times you have drunk alcohol or taken psychoactive substances, how many times did you do it...”). The consumption motives assessment investigated social (e.g., “because it helps you enjoy a party”), enhancement (e.g., “because you like the feeling”), conformity (e.g., “to fit in with a group you like”), and coping (e.g., “to cheer up when you are in a bad mood”) motives. The short form of this questionnaire showed comparable factor structure and similar concurrent validity to the original form (Cooper, 1994). This questionnaire also had a good internal consistency in the current sample (see Supplementary Table). All the participants included (66.8% women) were between 19 and 35 years old ($M = 20.79$; $SD = 2.83$) and were fluent French speakers. Although young adults are usually defined between 18 and 30 years old (e.g., Lipperman-Kreda, Paschall, Robert, & Morrison, 2018), we included all participants in this study⁴. Indeed, including all respondents allowed us to better cover substance use, psychological profiles, and personal characteristics.

²The response rate is line with the usual response rates observed at the University of Caen Normandie; e.g., 17.64% of students at the University of Caen Normandie answered a survey about blood donation in 2016; 11.41% answered a survey about healthcare access among students in 2018.

³Commission Nationale Informatique et Liberté.

⁴Analyses have been conducted in a subsample of participants between 18 and 30 years old and results were similar. We kept the whole sample for a better representation of the population.



Fig. 2. Subgroups of college students determined by cluster analysis according to impulsivity facets (urgency, lack of premeditation [Lprem], lack of perseverance [Lpers], sensation seeking [SS]) and consumption motives (social, enhancement, conformity, coping).

2.2.1. Substance use evaluation

Tobacco consumption was assessed using the Fagerström Test for Nicotine Dependence (FTND; [Heatherton, Kozlowski, Frecker, & Fagerström, 1991](#)), a 6-item test evaluating cigarette consumption and dependence (score range: 0–10). Scores below 4 indicate no dependence, scores between 4 and 5 indicate low dependence, whereas scores between 6 and 7 reflect moderate dependence, and scores higher than 7 high dependence.

Alcohol consumption was assessed by the Alcohol Use Disorders Identification Test (AUDIT; [Gache et al., 2005](#)), measuring the severity of alcohol consumption and alcohol use disorders using a 10-item Likert scale (score range: 0–40). AUDIT scores higher than 7 reflect hazardous consumption whereas scores beyond 19 indicate potential severe alcohol use disorder. Complementary items were also used to explore specific binge drinking habits, namely the consumption speed (i.e. number of alcohol doses consumed in one hour; an alcohol dose containing 10 gr of pure ethanol in France), the number of drunkenness episodes during the last 6 months, and the percentage of alcohol consumption episodes leading to drunkenness in the last six months. A binge drinking score ([Townshend & Duka, 2002, 2005](#)) was computed using the following formula: $[(4 \times \text{consumption speed}) + \text{drunkenness frequency} + (0.2 \times \text{drunkenness percentage})]$ to consider participants who drank heavily but irregularly.

Finally, cannabis, ecstasy, cocaine, and heroin consumptions were evaluated through the Cannabis Abuse Screening Test (CAST; [Legleye, Karila, Beck, & Reynaud, 2007](#)) adapted for each substance (score range: 0–24). This questionnaire evaluated consumption frequency and harmfulness (i.e., the prevalence of non-recreational use, memory disorders, inability to reduce or stop using the substance, and problems related to consumption). CAST scores beyond 2 reflect a harmful consumption. This test presented acceptable to very good internal consistencies (see Supplementary Table).

2.3. Data analysis

Descriptive analyses were first performed to explore the prevalence of substance use in the whole sample. A data clustering technique was then used to identify subgroups among this sample (including all people who had already use any substance). As previously performed ([Billieux et al., 2015; Hair, Black, Babin, & Anderson, 2010; Lannoy et al., 2017](#)), we conducted data grouping through a combination of hierarchical (using Ward's method with a squared Euclidean distance measure) and non-hierarchical (i.e., K-means analysis) methods. We designated the psychological variables selected for the cluster analysis according to established risk factors for substance use, i.e., impulsivity (urgency, lack of premeditation, lack of perseverance, sensation seeking) and motives (enhancement, social, conformity, coping). All the variables selected were Z-scored to ensure that they had the same metric

properties. To avoid multicollinearity ([Hair et al., 2010](#)), we also merged positive and negative urgency facets of impulsivity (correlation: $r = 0.52, p < 0.001$) into an “urgency” factor, in line with previous recommendations ([Billieux et al., 2015](#)). Finally, subgroups were compared based on external correlates (i.e., age, gender, academic years, academic attendance, anxiety, self-esteem, and substance use), post-hoc tests were corrected for multiple comparisons with Bonferroni procedure when appropriated. Regarding substance use comparisons, we explored: (1) the prevalence of each substance use in the different subgroups by binary evaluations (yes/no) and (2) the substance use in each cluster thanks to continuous scores. We investigated polysubstance consumption by computing the percentage of individuals using two substances (i.e., tobacco and cannabis) or all substances (i.e., tobacco, alcohol, cannabis, ecstasy, cocaine) in each cluster. As nearly all students drank alcohol, we did not explore polysubstance use in link with alcohol consumption.

3. Results

3.1. Descriptive analysis

Descriptive analyses showed that 42.1% of students smoke tobacco, 99.3% drink alcohol, 24.4% smoke cannabis, 4.9% consume ecstasy, 1.9% consume cocaine, and 0.2% consume heroin.

3.2. Cluster analysis

We performed cluster analysis on the whole sample, and results indicated an optimal four-factor solution ([Fig. 2](#)). The four clusters encompassed 33.9%, 31.8%, 21.1%, and 13.2% of the sample. Analyses of variance confirmed the differences between clusters regarding impulsivity and consumption motives. Comparisons based on external correlates strongly supported the reliability of these subgroups ([Table 1](#)). First, results showed that Cluster 1 was characterized by low impulsivity and consumption motives, Cluster 2 by high values of enhancement and social motives as well as average scores for sensation seeking and lack of perseverance, Cluster 3 by extreme conformity motives, elevated drinking motives, as well as intense urgency, and Cluster 4 by globally intense impulsivity (urgency, lack of perseverance, sensation seeking, lack of premeditation) and high motives (coping, enhancement, social). Second, the main finding observed in the cluster comparisons was that substance use significantly differed between clusters, with an increased prevalence in Cluster 4. The fourth cluster encompassed individuals presenting larger consumption of all substances, increased dangerousness, but also polysubstance use ([Fig. 3](#)). Although the sample showed a higher prevalence of women, gender ratio differed between clusters and was more balanced in Cluster 2 compared to the others. Regarding university years, results showed that

Table 1
Descriptive statistics and mean comparisons between clusters.

Variables	Cluster 1 (n = 930)	Cluster 2 (n = 872)	Cluster 3 (n = 362)	Cluster 4 (n = 577)	F/khi2	Comparisons
Cluster profiles [mean (SD)]						
Impulsivity						
Urgency	8.93 (2.15)	9.04 (1.83)	10.67 (2.11)	11.89 (1.89)	336.30*	C4 > C3 > C2 = C1
Lack of premeditation	6.71 (1.86)	6.94 (1.69)	7.09 (1.86)	9.66 (2.01)	355.51*	C4 > C3 > C2 = C1
Lack of perseverance	6.57 (2.14)	7.24 (2.23)	7.36 (2.41)	8.77 (2.51)	111.52*	C4 > C3 > C2 = C1
Sensation seeking	8.69 (2.76)	10.07 (2.73)	10.45 (2.84)	11.58 (2.56)	139.14*	C4 > C3 = C2 > C1
Consumption motives						
Social motives	5.23 (1.94)	10.59 (2.12)	11.57 (2.03)	10.18 (2.65)	1307.26*	C3 > C2 > C4 > C1
Enhancement motives	4.71 (1.71)	9.91 (2.29)	10.25 (2.53)	9.85 (2.74)	1108.75*	C3 = C4 = C2 > C1
Conformity motives	3.50 (1.10)	3.69 (1.07)	8.61 (2.20)	4.13 (1.48)	1361.13*	C3 > C4 > C2 < C1
Coping motives	3.89 (1.68)	4.82 (2.11)	7.61 (3.22)	7.88 (3.02)	452.71*	C4 = C3 > C2 > C1
External correlates [mean (SD)]						
Age	21.40 (3.49)	20.63 (2.39)	20.37 (2.28)	20.29 (2.38)	24.28*	C1 > C2 = C3 = C4
Age range	19–35	19–35	19–35	19–35		
Gender (% of women)	75.6	54.4	64.1	73.1	104.93*	C1 > C4 > C3 > C2
Academic year					98.08*	
First year (%)	30.4	29.2	13.7	26.7		
Second year (%)	28.3	36.3	12.5	22.9		
Third year (%)	36.4	32.2	14.4	17		
Fourth year (%)	38.2	33.5	14	14.3		
Fifth year (%)	45.1	32.2	12.4	10.3		
PhD students (%)	57.1	31.9	5.5	5.5		
Academic attendance (%)	88.19 (20.52)	83.68 (21.23)	82.70 (24.24)	78.52 (22.54)	24.39*	C1 > C2 = C3 > C4
Anxiety	45.16 (11.51)	42.88 (11.86)	49.51 (12.45)	50.62 (12.95)	59.20*	C4 = C3 > C1 > C2
Self-esteem	3.10 (1.26)	3.34 (1.18)	2.65 (1.30)	2.62 (1.37)	50.05*	C1 > C2 > C3 = C4
Tobacco consumption (FTND)	0.31 (1.11)	0.68 (1.46)	0.60 (1.41)	1.28 (1.99)	51.60*	C4 > C2 = C3 > C1
Alcohol consumption (AUDIT)	3.51 (2.84)	7.99 (4.95)	9.00 (5.60)	9.45 (6.02)	255.10*	C4 = C3 > C2 > C1
Binge drinking score	8.34 (9.53)	26.66 (20.36)	26.66 (21.84)	29.36 (23.08)	223.78*	C4 > C3 = C2 > C1
Cannabis consumption (CAST)	0.08 (0.61)	1.01 (2.59)	1.28 (3.44)	2.24 (4.18)	75.79*	C4 > C3 = C2 > C1
Ecstasy consumption (adapted CAST)	0.01 (0.12)	0.15 (0.89)	0.12 (0.77)	0.25 (1.10)	13.07*	C4 > C3 = C2 > C1
Cocaine consumption (adapted CAST)	0.00 (0.03)	0.03 (0.34)	0.01 (0.11)	0.14 (1.06)	9.32*	C4 > C3 = C2 = C1

Note. FTND = Fagerström Test for Nicotine Dependence; AUDIT = Alcohol Use Disorder Identification Test; CAST = Cannabis Abuse Screening Test. Cluster comparisons for the heroin CAST score were not performed as heroin users (n = 5) are regrouped in Cluster 4. Differences between clusters are shown by corrected Bonferroni post-hoc tests (excepted for gender and academic years). *p < 0.001.

participants in Cluster 1 were more advanced in their academic background (i.e., higher prevalence of master's and Ph.D. students) whereas participants of Cluster 4 were rather in their first university years. Finally, concerning psychopathological factors, findings highlighted that individuals of Cluster 1 reported higher self-esteem and individuals of Cluster 2 had lower anxiety. Members of Clusters 3 and 4 showed higher anxiety and lower self-esteem.

4. Discussion

The current study aimed to identify the existence of distinct psychological profiles of substance users among college students. The current findings support the existence of distinct psychological profiles and highlight four subgroups characterized by specific impulsivity facets and consumption motives. These profiles differ regarding the intensity and frequency of substance use, anxiety, and self-esteem. Moreover, while the current results highlight distinct profiles of substance users and identify those at risk for excessive use, these profiles were not characterized by a large variation regarding the type of substance consumed. This finding appears in contrast with previous studies investigating heterogeneity in social and environmental variables (Schilling et al., 2017; Stanley & Swaim, 2018). Regarding psychological factors, our results demonstrate specific characteristics related to multiple consumptions rather than profiles of alcohol or drug users.

First of all, consistent with previous studies showing that substance users have higher impulsivity and consumption motives than no-substance users (Charles et al., 2016; Dow & Kelly, 2013), the current research reveals that Cluster 1 is composed of young people with light substance use, as well as impulsivity and motives scores below the

mean of the whole sample. Indeed, these students report a preserved self-regulation (reduced impulsivity), a lack of desire to consume drugs, but also good self-esteem and low anxiety. Moreover, this cluster is characterized by a stronger prevalence of women and older students, which widely supports the influence of gender and age on substance use (Degenhardt et al., 2008).

Regarding the subgroups of people using substances, Cluster 2 is characterized by important enhancement and social motives. Accordingly, we postulate that this cluster includes people with a classical "student profile" who use substances for pleasure and sensations in social and festive contexts. Indeed, it has been shown that emerging adulthood is associated with increased autonomy and freedom, leading to new experiences such as substance use (see Stone, Becker, Huber, & Catalano, 2012, for a review). The university environment can further facilitate this freedom and can lead young people to explore new experiences. This proposal is reinforced by the apparent preservation of psychological well-being in this subgroup (good self-esteem, academic attendance, and low anxiety), supporting that substance use would be mainly contextual and not related to psychological difficulties. Therefore, we expect that most students in this cluster would evolve towards a low-risk profile with studies' progress or at the beginning of their professional life. This assumption is in line with existing longitudinal studies showing that leaving University is related to a decrease of substance use in people with low-risk profiles (e.g., Gómez et al., 2017). However, it is worth noting that these individuals also report high scores for the use of all substances and elevated binge drinking. These consumption habits may have deleterious consequences at short-term but also in the long-term (e.g., cognitive and cerebral impairments; Mahmood et al., 2013; McNamee et al., 2008). Prevention




Fig. 3. Substance use in the four clusters. (a) Represents the prevalence of substance use in each cluster (i.e., tobacco, alcohol, cannabis, ecstasy, cocaine, heroin), the percentage of individuals consuming both tobacco and cannabis or all substances. (b) Illustrates substance use in each cluster, in reference to validated tools (Fagerström Test for Nicotine Dependence [FTND], Alcohol Use Disorder Identification Test [AUDIT], Cannabis Abuse Screening Test [CAST] for cannabis, ecstasy, cocaine, and heroin consumption). Scores presented in the Y axis are standardized in order to reliably compare the different tests.

actions modifying the potential misperception that substance use is normative and informing about the negative consequences (Botvin & Griffin, 2007; Onrust, van der Heijden, Zschämisch, & Speetjens, 2018) should thus be carried out among individuals in this cluster.

Cluster 3 is mainly defined by important conformity, together with enhancement, social, and coping motives, as well as high urgency scores. Other impulsivity facets, although slightly more elevated than in clusters 1 and 2, are close to the average of the whole sample. This cluster characterizes a part of substance users who particularly use alcohol and drug to avoid being excluded by others (Studer et al., 2016). Subsequent analyses also showed that college students in this subgroup have important levels of anxiety and low self-esteem. Therefore, this third subgroup is comparable to the second one in terms of substance use but present lower psychological well-being. Due to its mean impulsivity scores (lack of premeditation, lack of perseverance, sensation seeking), this cluster contrasts with the current literature emphasizing the crucial role of impulsivity in substance use (e.g., Charles et al., 2016; Loree et al., 2015). These results show that people with low personal risk factors to use substances may still have significant substance consumption and poor well-being. This finding is, however, in accordance with previous studies showing the unique predictive value of consumption motives (Dow & Kelly, 2013) and underlines the dangerousness related to negative reinforcement motives (e.g., Cooper, 1994; Dow & Kelly, 2013; Mezquita et al., 2011). It constitutes a main insight of the present study as it highlights a currently under-reported

category of college students. Indeed, in this subgroup, we observe a relationship between substance use, peer pressure (conformity), and stress (coping), possibly related to the university environment (e.g., the burden to success in a high social environment). This subgroup emphasizes the need for prevention programs improving the ability to refuse drugs and to resist peer influence (Botvin & Griffin, 2007; Helmer, Muellmann, Zeeb, & Pischke, 2016).

Finally, Cluster 4 encompasses another part of substance users and show a “classical profile” of heavy consumers, characterized by elevated impulsivity and consumption motives (coping, enhancement, social), low self-esteem, low academic attendance, and high anxiety. This psychological profile is in line with previous studies, showing a strong influence of impulsivity and emotional motives in substance use (e.g., Barahmand, Khazaei, & Hashjin, 2016; Cooper, Wood, Orcutt, & Albino, 2003; Lammers et al., 2013) as well as a significant relationship with anxiety (e.g., Tavalacci et al., 2013). Accordingly, we postulate that participants in this subgroup use substances to increase positive affects (enhancement, social, urgency, sensation seeking) or decrease negative ones (coping, urgency), while reporting low perseverance and few considerations of the long-term consequences. The hazardousness of this profile is observed through elevated substance use and higher prevalence of polysubstance use. Previous studies showed that polysubstance use is related to increased mental illness and violent/high-risk behaviors compared to unique substance use (Connora et al., 2014; Morley, Lynskey, Moran, Borschmann, & Winstock, 2015). This result

thus reinforces the negative health and social outcomes of this cluster. The possible consequences of such a psychological profile of substance users should be underlined. As students in this subgroup present the typical risk profile for substance abuse and dependence, they are at risk of evolving towards severe substance use disorders after their university years. Prevention and intervention programs are critical for this subgroup and might focus on improving personal skills (e.g., emotion regulation, perseverance, premeditation), cognitive restructuring (Botvin & Griffin, 2007; Chambers et al., 2016), or cognitive stimulation (Sampedro-Piquero et al., 2018) to avoid such transition towards established addictive disorders.

Importantly, Clusters 3 and 4 are characterized by psychological problems (high anxiety and low self-esteem), which may no longer reflect recreational substance use. The association between substance use and anxiety is indeed highly prevalent in clinical populations and can be explained by the self-medication hypothesis (for sedatives; alcohol, cannabis) or the neurotoxic effect of substance use (for stimulants; tobacco, cocaine, heroin) (Vorspan, Mehtelli, Dupuy, Bloch, & Lépine, 2015). Here, we observe this relationship for individuals presenting the consumption of both sedative and stimulant substances, with a higher prevalence of sedative substances use (i.e., alcohol and cannabis). According to this observation and the age of our sample, we hypothesize that some participants in these clusters use substances as self-medication, which is also consistent with the coping motives and urgency facet observed in these clusters. Moreover, these results can also be related to a recent study showing subgroups characterized by either high self-esteem without psychological problems or low self-esteem with psychological problems (Gierski et al., 2020). Altogether, these results highlight the significant role of anxiety and self-esteem to characterize these subgroups.

Some limitations should be acknowledged. First, although comparable with previous studies, the response rate in the current research is moderate (13.87%). Second, whereas the consumption motives questionnaire used in this study is in line with previous studies (Cooper et al., 2016) and has good reliability in the current sample, future research should reinforce our results by using substance-specific motives scales. In the same vein, self-esteem was assessed by a single item with a likert scale (Robins et al., 2001). However, given the important role of self-esteem in these findings, this concept should be further explored. Overall, this exploration of distinct psychological profiles among substance users has to be extended through experimental and longitudinal data. It would also be interesting to consider other substances use (e.g., nicotine-cigarette, methamphetamine, inhalants). Finally, although these results are consistent with previous French studies (e.g., Gierski et al., 2020; Tavalacci et al., 2013), they should be replicated by targeting adults with various ages, socio-economic status, and from various countries.

As a whole, the current results show distinct psychological profiles of college students and highlight several patterns of combined impulsivity and motivations related to excessive substance use. The associations found between impulsivity facets (urgency, lack of premeditation, sensation seeking) and drinking motives (coping, social, enhancement) in clusters 3 and 4 extend previous results, showing the interaction between these variables to explain excessive/problematic alcohol use (Adams et al., 2012; Jones et al., 2014). In addition, our results highlight the joint role of conformity motives and urgency in Cluster 3. In particular, by indicating profiles of recreational (enhancement and social motives), peer pressure-sensitive (high conformity motives), and hazardous (high impulsivity and consumption motives) users, these findings are in line with previous ones (Lannoy et al., 2017). This study also goes beyond the widely described role of impulsivity by showing that people with good self-regulation also report substance use. Indeed, Cluster 3 highlights that people with good executive abilities may exhibit harmful substance use, mainly explained by a need to comply. These results thus have strong implications to target substance users and implement appropriate prevention and

intervention actions. They suggest that prevention of substance use should be adapted according to the psychological characteristics of each profile rather than being conducted in association with the substance consumed.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This research was supported by a grant from the Fondation pour la Recherche en Alcoologie (FRA, France), Séverine Lannoy was funded by the Belgian American Educational Foundation (BAEF, Belgium, United States of America), and Pierre Maurage by the National Fund for Scientific Research (FRS-FNRS, Belgium), but these funds did not exert any editorial direction or censorship on any part of this article.

Contributors

HB, JM, PL, and LR were responsible for the research implementation and acquisition. HB, JM, PM, and SL design the current study and all authors participated in its construction. SL conducted literature searches and provided summaries of previous research studies. SL and PM performed the statistical analysis. SL wrote the first draft of the manuscript and all authors contributed to and have approved the final manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.addbeh.2020.106477>.

References

- Adams, Z. W., Kaiser, A. J., Lynam, D. R., Charnigo, R. J., & Milich, R. (2012). Drinking motives as mediators of the impulsivity-substance use relation: Pathways for negative urgency, lack of premeditation, and sensation seeking. *Addictive Behaviors*, *37*, 848–855. <https://doi.org/10.1016/j.addbeh.2012.03.016>.
- Assanangkornchai, S., Li, J., McNeil, E., & Saingam, D. (2018). Clusters of alcohol and drug use and other health-risk behaviors among Thai secondary school students: A latent class analysis. *BMC Public Health*, *18*, 1272. <https://doi.org/10.1186/s12889-018-6205-z>.
- Ayala, E. E., Roseman, D., Winseman, J. S., & Mason, H. R. C. (2017). Prevalence, perceptions, and consequences of substance use in medical students. *Medical Education*, *22*, 1392824. <https://doi.org/10.1080/10872981.2017.1392824>.
- Barahmand, U., Khazaei, A., & Hashjin, G. S. (2016). Emotion dysregulation mediates between childhood emotional abuse and motives for substance use. *Archives of Psychiatric Nursing*, *30*, 653–659. <https://doi.org/10.1016/j.apnu.2016.02.007>.
- Billieux, J., Rochat, L., Ceschi, G., Carré, A., Offerlin-Meyer, I., Defeldre, A.-C., ... Van der Linden, M. (2012). Validation of a short French version of the UPPS-P impulsive behavior scale. *Comprehensive Psychiatry*, *53*, 609–615. <https://doi.org/10.1016/j.comppsych.2011.09.001>.
- Billieux, J., Thorens, G., Khazaal, Y., Zullino, D., Achab, S., & Van der Linden, M. (2015). Problematic involvement in online games: A cluster analytic approach. *Computers in Human Behavior*, *43*, 242–250. <https://doi.org/10.1016/j.chb.2014.10.055>.
- Blank, M.-L., Connor, J., Gray, A., & Tustin, K. (2016). Alcohol use, mental well-being, self-esteem and general self-efficacy among final-year university students. *Social Psychiatry and Psychiatric Epidemiology*, *51*, 431–441. <https://doi.org/10.1007/s00127-016-1183-x>.
- Botvin, G. J., & Griffin, K. W. (2007). School-based programmes to prevent alcohol, tobacco and other drug use. *International Review of Psychiatry*, *19*, 607–615. <https://doi.org/10.1080/09540260701797753>.
- Bruchon-Schweitzer, M., & Paulhan, I. (1993). *Adaptation francophone de l'inventaire d'anxiété Trait-État (Forme Y) de Spielberger*. Paris: Editions du Centre de Psychologie Appliquée.
- Center for Behavioral Health Statistics and Quality. (2015). Behavioral health trends in the United States: Results from the 2014 National Survey on Drug Use and Health (HHS Publication No. SMA 15-4927, NSDUH Series H-50). Retrieved from <http://www.samhsa.gov/data/>.
- Chambers, J. E., Brooks, A. C., Medvin, R., Metzger, D. S., Lauby, J., Carpenedo, C. M., ... Kirby, K. C. (2016). Examining multi-session brief intervention for substance use in

- primary care: Research methods of a randomized controlled trial. *Addiction Science & Clinical Practice*, 11. <https://doi.org/10.1186/s13722-016-0057-6>.
- Charles, N. E., Ryan, S. R., Bray, B. C., Mathias, C. W., Acheson, A., & Dougherty, D. M. (2016). Altered developmental trajectories for impulsivity and sensation seeking among adolescent substance users. *Addictive Behaviors*, 60, 235–241. <https://doi.org/10.1016/j.addbeh.2016.04.016>.
- Connora, J. P., Gullo, M. J., White, A., & Kelly, A. B. (2014). Polysubstance use: diagnostic challenges, patterns of use and health. *Current Opinion in Psychiatry*, 27, 269–275. <https://doi.org/10.1097/YCO.0000000000000069>.
- Cooper, M. L. (1994). Motivations for alcohol use among adolescents: Development and validation of a four-factor model. *Psychological Assessment*, 6, 117–128. <https://doi.org/10.1037/1040-3590.6.2.117>.
- Cooper, M. L., Kuntsche, E., Levitt, A., Barber, L. L., & Wolf, S. (2016). *Motivational models of substance use: A review of theory and research on motives for using alcohol, marijuana, and tobacco*. The Oxford handbook of substance use disorders. Oxford University Press 1–53.
- Cooper, M. L., Wood, P. K., Orcutt, H. K., & Albino, A. (2003). Personality and the predisposition to engage in risky or problem behaviors during adolescence. *Journal of Personality and Social Psychology*, 84, 390–410. <https://doi.org/10.1037/0022-3514.84.2.390>.
- Degenhardt, L., Chiu, W.-T., Sampson, N., Kessler, R. C., Anthony, J. C., Angermeyer, M., ... Wells, J. E. (2008). Toward a global view of alcohol, tobacco, cannabis, and cocaine use: Findings from the WHO World Mental Health Surveys. *PLoS Medicine*, 5, e141. <https://doi.org/10.1371/journal.pmed.0050141>.
- Dow, S. J., & Kelly, J. F. (2013). Listening to youth: Adolescents' reasons for substance use as a unique predictor of treatment response and outcome. *Psychology of Addictive Behaviors*, 27, 1122–1131. <https://doi.org/10.1037/a0031065>.
- Ehret, P. J., Ghaidarov, T. M., & LaBrie, J. W. (2013). Can you say no? Examining the relationship between drinking refusal self-efficacy and protective behavioral strategy use on alcohol outcomes. *Addictive Behaviors*, 38, 1898–1904. <https://doi.org/10.1016/j.addbeh.2012.12.022>.
- Evans-Polce, R., Lanza, S., & Maggs, J. (2016). Heterogeneity of alcohol, tobacco, and other substance use behaviors in U.S. college students: A latent class analysis. *Addictive Behaviors*, 53, 80–85. <https://doi.org/10.1016/j.addbeh.2015.10.010>.
- Fitzgerald, A., Mac Giollaibhui, N., Dolphin, L., Whelan, R., & Dooley, B. (2018). Dissociable psychosocial profiles of adolescent substance users. *PLoS ONE*, 13, e0202498. <https://doi.org/10.1371/journal.pone.0202498>.
- Gache, P., Michaud, P., Landry, U., Accietto, C., Arfaoui, S., Wenger, O., & Daeppen, J.-B. (2005). The alcohol use disorders identification test (AUDIT) as a screening tool for excessive drinking in primary care: Reliability and validity of a French version. *Alcoholism, Clinical and Experimental Research*, 29, 2001–2007. <https://doi.org/10.1097/01.alc.0000187034.58955.64>.
- Gierski, F., Benzerouk, F., De Wever, E., Duka, T., Kaladjian, A., Quaglini, V., & Naassila, M. (2017). Cloninger's temperament and character dimensions of personality and binge drinking among college students. *Alcoholism, Clinical and Experimental Research*, 41, 1970–1979. <https://doi.org/10.1111/acer.13497>.
- Gierski, F., De Wever, E., Benzerouk, F., Lannoy, S., Kaladjian, A., Naassila, M., & Quaglini, V. (2020). Disentangling the relationship between self-esteem and problematic alcohol use among college students: Evidence from a cluster analytic approach. *Alcohol and Alcoholism*, 55, 196–203. <https://doi.org/10.1093/alcalc/azg097>.
- Gómez, P., Moure-Rodríguez, L., López-Caneda, E., Rial, A., Cadaveira, F., & Caamaño-Isorna, F. (2017). Patterns of alcohol consumption in Spanish university alumni: Nine years of follow-up. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.00756>.
- Grant, V. V., Stewart, S. H., O'Connor, R. M., Blackwell, E., & Conrod, P. J. (2007). Psychometric evaluation of the five-factor Modified Drinking Motives Questionnaire—Revised in undergraduates. *Addictive Behaviors*, 32, 2611–2632. <https://doi.org/10.1016/j.addbeh.2007.07.004>.
- Hair, J. F., Black, W. C., Babin, B. J., & Anderson, R. E. (2010). *Multivariate data analysis: A global perspective*. London: Pearson Education.
- Heatherton, T. D., Kozlowski, L. T., Frecker, R. C., & Fagerström, K.-O. (1991). The fagerström Test for nicotine dependence: A revision of the Fagerström Tolerance Questionnaire. *British Journal of Addiction*, 86, 1119–1127.
- Hecimovic, K., Barrett, S. P., Darredeau, C., & Stewart, S. H. (2014). Cannabis use motives and personality risk factors. *Addictive Behaviors*, 39, 729–732. <https://doi.org/10.1016/j.addbeh.2013.11.025>.
- Helmer, S. M., Muellmann, S., Zeeb, H., & Pischke, C. R. (2016). Development and evaluation of the efficacy of a web-based 'social norms'-intervention for the prevention and reduction of substance use in a cluster-controlled trial conducted at eight German universities. *BMC Public Health*, 16, 252. <https://doi.org/10.1186/s12889-016-2898-z>.
- Hides, L., Lubman, D. I., Cosgrave, E. M., Buckley, J. A., Killackey, E., & Yung, A. R. (2008). Motives for substance use among young people seeking mental health treatment. *Early Intervention in Psychiatry*, 2, 188–194. <https://doi.org/10.1111/j.1751-7893.2008.00076.x>.
- Jones, K. A., Chryssanthakis, A., & Groom, M. J. (2014). Impulsivity and drinking motives predict problem behaviours relating to alcohol use in University students. *Addictive Behaviors*, 39, 289–296. <https://doi.org/10.1016/j.addbeh.2013.10.024>.
- Kuntsche, E., & Kuntsche, S. (2009). Development and validation of the drinking motive questionnaire revised short form (DMQ-R SF). *Journal of Clinical Child and Adolescent Psychology*, 38, 899–908. <https://doi.org/10.1080/15374410903258967>.
- Lammers, J., Kuntsche, E., Engels, R. C. M. E., Wiers, R. W., & Kleinjan, M. (2013). Medial relations of substance use risk profiles, alcohol-related outcomes, and drinking motives among young adolescents in the Netherlands. *Drug and Alcohol Dependence*, 133, 571–579. <https://doi.org/10.1016/j.drugalcdep.2013.07.030>.
- Lannoy, S., Billieux, J., Poncin, M., & Muraire, P. (2017). Binge at the campus: Motivations and impulsivity influence binge drinking profiles in university students. *Psychiatry Research*, 250, 146–154. <https://doi.org/10.1016/j.psychres.2017.01.068>.
- Lee, C. A., Derefiniko, K. J., Davis, H. A., Milich, R., & Lynam, D. R. (2017). Cross-lagged relations between motives and substance use: Can use strengthen your motivation over time? *Drug and Alcohol Dependence*, 178, 544–550. <https://doi.org/10.1016/j.drugalcdep.2017.05.027>.
- Legleye, S., Karila, L., Beck, F., & Reynaud, M. (2007). Validation of the CAST, a general population Cannabis Abuse Screening Test. *Journal of Substance Misuse*, 12, 233–242. <https://doi.org/10.1080/14659890701476532>.
- Lippman-Kreda, S., Paschall, M. J., Robert, S., & Morrison, C. N. (2018). Places and social contexts associated with simultaneous use of alcohol, tobacco and marijuana among young adults: Social contexts of simultaneous use. *Drug and Alcohol Review*, 37, 188–195. <https://doi.org/10.1111/dar.12537>.
- Loree, A. M., Lundahl, L. H., & Ledgerwood, D. M. (2015). Impulsivity as a predictor of treatment outcome in substance use disorders: Review and synthesis: Impulsivity in substance use treatment. *Drug and Alcohol Review*, 34, 119–134. <https://doi.org/10.1111/dar.12132>.
- Magallón-Neri, E., Díaz, R., Fornis, M., Goti, J., & Castro-Fornieles, J. (2015). Personality psychopathology, drug use and psychological symptoms in adolescents with substance use disorders and community controls. *PeerJ*, 3, e992. <https://doi.org/10.7717/peerj.992>.
- Mahmood, O. M., Goldenberg, D., Thayer, R., Migliorini, R., Simmons, A. N., & Tapert, S. F. (2013). Adolescents' fMRI activation to a response inhibition task predicts future substance use. *Addictive Behaviors*, 38, 1435–1441. <https://doi.org/10.1016/j.addbeh.2012.07.012>.
- Malmberg, M., Kleinjan, M., Vermulst, A. A., Overbeek, G., Monshouwer, K., Lammers, J., & Engels, R. C. M. E. (2012). Do substance use risk personality dimensions predict the onset of substance use in early adolescence? A variable- and Person-centered approach. *Journal of Youth and Adolescence*, 41, 1512–1525. <https://doi.org/10.1007/s10964-012-9775-6>.
- Martinez-Loredo, V., Fernandez-Hermida, J. R., De La Torre-Luque, A., & Fernandez-Artamendi, S. (2018). Trajectories of impulsivity by sex predict substance use and heavy drinking. *Addictive Behaviors*, 85, 164–172. <https://doi.org/10.1016/j.addbeh.2018.06.011>.
- McCabe, S. E., Boyd, C. J., Couper, M. P., Crawford, S., & D'arcy, H. (2002). Mode effects for collecting alcohol and other drug use data: Web and U.S. Mail. *Journal of Studies on Alcohol and Drugs*, 63, 755–761.
- McNamee, R. L., Dunfee, K. L., Luna, B., Clark, D. B., Eddy, W. F., & Tarter, R. E. (2008). Brain activation, response inhibition, and increased risk for substance use disorder. *Alcoholism, Clinical and Experimental Research*, 32, 405–413. <https://doi.org/10.1111/j.1530-0277.2007.00604.x>.
- Mezquita, L., Stewart, S. H., Ibáñez, M. I., Ruipérez, M. A., Villa, H., Moya, J., & Ortet, G. (2011). Drinking motives in clinical and general populations. *European Addiction Research*, 17, 250–261. <https://doi.org/10.1159/000328510>.
- Morley, K. I., Lynskey, M. T., Moran, P., Borschmann, O., & Winstock, A. R. (2015). Polysubstance use, mental health and high-risk behaviours: Results from the 2012 Global Drug Survey. *Drug and Alcohol Review*, 34, 427–437. <https://doi.org/10.1111/dar.12263>.
- Neighbors, C., Dillard, A. J., Lewis, M. A., Bergstrom, R. L., & Neil, T. A. (2006). Normative misperceptions and temporal precedence of perceived norms and drinking. *Journal of Studies on Alcohol*, 67, 290–299.
- Observatoire Européen des Drogues et des Toxicomanie (2017) Rapport européen sur les drogues : Tendances et évolutions.
- Onrust, S. A., van der Heijden, A., Zschämisch, A. L., & Speetjens, P. A. M. (2018). Effectiveness of fresh start: A randomized study of a school-based program to retain a negative attitude toward substance use in secondary school freshmen. *Substance Use & Misuse*, 53, 921–930. <https://doi.org/10.1080/10826084.2017.1385082>.
- Patrick, M. E., Fairlie, A. M., & Lee, C. M. (2018). Motives for simultaneous alcohol and marijuana use among young adults. *Addictive Behaviors*, 76, 363–369. <https://doi.org/10.1016/j.addbeh.2017.08.027>.
- Pedersen, M. U., Thomsen, R. K., Pedersen, M. M., & Hesse, M. (2017). Mapping risk factors for substance use: Introducing the YouthMap12. *Addictive Behaviors*, 65, 40–50. <https://doi.org/10.1016/j.addbeh.2016.09.005>.
- Robins, R. W., Hendin, H. M., & Trzesniewski, K. H. (2001). Measuring global self-esteem: Construct validation of a single-item measure and the Rosenberg self-esteem scale. *Personality and Social Psychology Bulletin*, 27, 151–161. <https://doi.org/10.1177/0146167201272002>.
- Thomsen, R. K., Callesen, M. B., Hesse, M., Kvamme, T. L., Pedersen, M. M., Pedersen, M. U., & Voon, V. (2018). Impulsivity traits and addiction-related behaviors in youth. *Journal of Behavioral Addictions*, 7, 317–330. <https://doi.org/10.1556/2006.7.2018.22>.
- Sampedro-Piquero, P., Ladrón de Guevara-Miranda, D., Pavón, F. J., Serrano, A., Suárez, J., Rodríguez de Fonseca, F., ... Castilla-Ortega, E. (2018). Neuroplastic and cognitive impairment in substance use disorders: A therapeutic potential of cognitive stimulation. *Neuroscience & Biobehavioral Reviews*, 106, 23–48. <https://doi.org/10.1016/j.neubiorev.2018.11.015>.
- Schilling, L., Zeeb, H., Pischke, C., Helmer, S., Schmidt-Pokrzywniak, A., Reintjes, R., ... Schneider, S. (2017). Licit and illicit substance use patterns among university students in Germany using cluster analysis. *Substance Abuse Treatment, Prevention, and Policy*, 12, 44. <https://doi.org/10.1186/s13011-017-0128-z>.
- Shin, S. H., Hong, H. G., & Jeon, S.-M. (2012). Personality and alcohol use: The role of impulsivity. *Addictive Behaviors*, 37, 102–107. <https://doi.org/10.1016/j.addbeh.2011.09.006>.
- Skalisky, J., Wielgus, M. D., Aldrich, J. T., & Mezulis, A. H. (2019). Motives for and impairment associated with alcohol and marijuana use among college students.

- Addictive Behaviors*, 88, 137–143. <https://doi.org/10.1016/j.addbeh.2018.08.028>.
- Skidmore, C. R., Kaufman, E. A., & Crowell, S. E. (2016). Substance use among college students. *Child and Adolescent Psychiatric Clinics of North America*, 25, 735–753. <https://doi.org/10.1016/j.chc.2016.06.004>.
- Stone, A. L., Becker, L. G., Huber, A. M., & Catalano, R. F. (2012). Review of risk and protective factors of substance use and problem use in emerging adulthood. *Addictive Behaviors*, 3, 747–775. <https://doi.org/10.1016/j.addbeh.2012.02.014>.
- Stanley, L. R., & Swaim, R. C. (2018). Latent classes of substance use among American Indian and White students living on or near reservations, 2009–2013. *Public Health Reports*, 133, 432–441. <https://doi.org/10.1177/0033354918772053>.
- Studer, J., Baggio, S., Grazioli, V. S., Mohler-Kuo, M., Daepfen, J.-B., & Gmel, G. (2016). Risky substance use and peer pressure in Swiss young men: Test of moderation effects. *Drug and Alcohol Dependence*, 168, 89–98. <https://doi.org/10.1016/j.drugalcdep.2016.08.633>.
- Tavolacci, M. P., Ladner, J., Grigioni, S., Richard, L., Villet, H., & Dechelotte, P. (2013). Prevalence and association of perceived stress, substance use and behavioral addictions: A cross-sectional study among university students in France, 2009–2011. *BMC Public Health*, 13. <https://doi.org/10.1186/1471-2458-13-724>.
- Townshend, J. M., & Duka, T. (2002). Patterns of alcohol drinking in a population of young social drinkers: A comparison of questionnaire and diary measures. *Alcohol and Alcoholism*, 37, 187–192.
- Townshend, J. M., & Duka, T. (2005). Binge drinking, cognitive performance and mood in a population of young social drinkers. *Alcoholism, Clinical and Experimental Research*, 29, 317–325. <https://doi.org/10.1097/01.ALC.0000156453.05028.F5>.
- Vorspan, F., Mehtelli, W., Dupuy, G., Bloch, V., & Lépine, J.-P. (2015). Anxiety and substance use disorders: Co-occurrence and clinical issues. *Current Psychiatry Reports*, 17, 1–7. <https://doi.org/10.1007/s11920-014-0544-y>.
- Walther, B., Morgenstern, M., & Hanewinkel, R. (2012). Co-occurrence of addictive behaviours: Personality factors related to substance use, gambling and computer gaming. *European Addiction Research*, 18, 167–174. <https://doi.org/10.1159/000335662>.
- Whiteside, S. P., & Lynam, D. R. (2001). The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, 30, 669–689. <https://doi.org/10.1002/ab.21474>.
- Wong, C. F., Silva, K., Kecojevic, A., Schrage, S. M., Bloom, J. J., Iverson, E., & Lankenau, S. E. (2013). Coping and emotion regulation profiles as predictors of nonmedical prescription drug and illicit drug use among high-risk young adults. *Drug and Alcohol Dependence*, 132, 165–171. <https://doi.org/10.1016/j.drugalcdep.2013.01.02>.